

## **Terms and Conditions**

The Library provides access to digitized documents strictly for noncommercial educational, research and private purposes and makes no warranty with regard to their use for other purposes. Some of our collections are protected by copyright. Publication and/or broadcast in any form (including electronic) requires prior written permission from the Library.

Each copy of any part of this document must contain there Terms and Conditions. With the usage of the library's online system to access or download a digitized document you accept there Terms and Conditions.

Reproductions of material on the web site may not be made for or donated to other repositories, nor may be further reproduced without written permission from the Library

For reproduction requests and permissions, please contact us. If citing materials, please give proper attribution of the source.

### Imprint:

Director: Mag. Renate Plöchl

Deputy director: Mag. Julian Sagmeister

Owner of medium: Oberösterreichische Landesbibliothek

Publisher: Oberösterreichische Landesbibliothek, 4021 Linz, Schillerplatz 2

### Contact:

Email: [landesbibliothek\(at\)ooe.gv.at](mailto:landesbibliothek(at)ooe.gv.at)

Telephone: +43(732) 7720-53100

tionem subministrarem: & quia incertum est, correxerint Arabes in Ptolemæicis istis aliqua, an omnis vetitionum diversitas à transcriptorum incuria sit orta. Pauca sunt, quibus ego manum admovi, vel in libro de stella Serpentarij vel aliàs, quas caractere antiquo, monendi causa, expressi.

**DENOMINATIO STELLARUM.** Longitudo G.M.S. Latitudo G.M. **DENOMINATIO STELLARUM.** Longitudo G.M.S. Latitudo G.M.

In URS Aminore informis Clavij (signo falso) est 8 Tychonis.  
In URS A majore 39.40.41, in meo Exemplari sunt magn.  
non 3, sed 5, lat. 34.59', 21.38', 20.44', & Ptol. 8 va in-  
formatarum inter eas esse videtur, quia Trapezuntius pro o  
habet o Q.

In CEPHEO, informis pr. Tiaram 4.30 V 64.0 B 5  
In BOOTE, 10 Ptol. in venabulo 28.45 H 45.45 4  
In ENGONASI in coxa sinist. pr. 1.20 H 55.55 5  
Sequens 2.30 H 58.15 5  
Informis dextri brachij Australior 24.39 M 57.30 5  
In OLORE in dextro genu 3.0 H 63.20 ne  
In PERSEO, informis à sinistro genu ad ortum 2.10 II 17.50 B 5

In OPHIUCHO & desunt multæ Ptolemæicæ, & suspectæ sunt aliquæ, quas Clavius ex Pifero sumpsit, qui aliàs merum Catalogum Tychonicum complexus est: cum tamen desint in meo exemplari. Quarum altitudines & culminationes cum instrumentis ceperim; Catalogum ex libro meo de stella nova transcribam, nihil præjudicans observatori diligentiori.

In recta trium ante pedē præ. bor. 25.42 M 9.27 B 5  
Media 24.47 M 6.42 5  
Infima 24.5 M 3.45 5  
Quæ præcedit genu anterius 1.3 H 13.34 5  
Duarum inter pedes superior 6.51 H 11.50 6  
Inferior 5.49 H 4.38 6  
Trium in crure anteriori borea 2.47 H 5.42 4  
o. Media 2.7 H 3.11 4  
p. Australissima 1.45 H 1.38 4  
q. In calcaneo pedis hujus 4.4 0.26 4  
In femore sequenti 15.39 17.28 5  
m. In cubito sequenti 19.2 H 14.57 4  
a. In pedis sequentis Tibia 15.42 H 1.57 B 4  
a. In digito pedis 14.7 H 3.37 A 4  
In dorso pedis 14.47 H 1.27 5  
b. In Plantâ, clara 15.47 H 1.43 3  
c. In vola 16.37 H 0.59 4  
d. e. In calce seu talo 18.5 H 0.57 A 4  
Sequens pedem hunc 20.4 H 1.22 B 4  
Infor. inter hum. Oph. & Herc. 26.27 M 35.0 B 4  
bb. Ad humerum Oph. in Rhomboborealissima 25.7 H 28.0 4  
cc. Mediarum præcedens 24.57 H 26.40 4  
dd. Sequens 26.50 H 26.28 4  
ee. Infima 25.40 H 24.45 4  
Stella nova anni 1604 17.40 H 1.56  
Supra informatas solitaria 27.50 H 32.45 B 4  
In SERPENTE, Quæ supra duplicem, octava Ptolemæo 16.33 M 26.36 B 4  
In crista 18.9 M 37.15 5  
In spira, apex Isofelis 13 Ptol. 15.0 H 10.25 4  
Basis præcedens 14 Ptol. 19.23 H 8.4 4  
Sequens 15 Ptol. 20.9 H 10.23 4  
Supra caudam duarum superior 29.49 H 26.38 6  
Inferior 1.39 P 23.28 6  
Infra caudam clara 3.0 P 15.49 4  
Trium inter ultimas caudæ ima 2.0 P 21.29 6  
Media 4.6 P 22.42 6  
Postrema 6.40 P 24.52 B 6

Proximè infra ultimam 10.22 P 25.2 B 6  
Sequens manum Ophiuchi 29.45 M 16.0 5  
In ANDROMEDA In Syrmatæ 4.0 H 34.40 5  
Australina [duarum bor. 5.30 H 32.40 5  
Inform. præced. tres in dextra manu 3.0 V 44.0 B 3  
In ARIETE. Trium infor. borea 12.20 H 12.44 B 5  
In extremo pede posteriori 6.0 H 5.10 A 4

In TAURO. Observatio de Plejade.  
Commendat eam creber appulsus Lunæ, & usus in obs. antiquissima magni momenti. Igitur Tycho solas 4 enumerat: & tamen primam earum non invenio in Observationibus. Etiamq; meum Msc. pro 23.13', habet 23.50', quod Bayeri disunctio probat. Videtur indulta Ptolemæo longitudo tam parva, ut qui etiam ad 1.30', dislocat terminos Plejadis, & sic usurpat in obs. Timocharidis: cum tamen hæc non ferat tantam. Idem Ptol. exorsus à quadrilatero, deficit in ejus enumeratione, quarto adiciens unam extrâ. Sunt tamen 7, quas Galilæus in Nuncio exprimit. At meæ disputationes diversissimorum temporum, tribus Tychonicis innixæ, consentiunt satis propinquè in hos numeros; ac si sic in Ptol. sit legendum.

τῆς Πλειάδος τὸ βόρειον πέρασ τῆς ἠγεμένης πλευρᾶς	24.5 H	4.21
τὸ νότιον πέρασ τῆς ἠγεμένης πλευρᾶς	23.50 H	4.13
τῆς ἐπομένης τὸ βόρειον πέρασ	24.24 H	4.3
τὸ νοτιώτατον πέρασ τῆς πλευρᾶς vel πλειάδος	24.4 H	3.55
ὁ ἐκτὸς τῆς πλειάδος, καὶ μικρὸς ἀπ' ἀριστερῶν. Ita deessent	24.7 H	4.25
ὁ ἠγεμενὸς αὐτῆς συχνὰ ἐκλείπων	24.0 H	4.26
ὁ ἐπομενὸς τῆς πλειάδος	24.47 H	3.55

Ac si ἡ πλειάσ sit Ratis, & contineatur solo quadrilatero. Et si etiam in ultima, observata altitudo, calculum superat. Omnino omiffam esse unam, Summa 32 arguit, sunt enim 33. Et Mag. primæ refertur una, tertiæ 6, cum clara Plejadis non sit minor eoq; septima, tertiæ Mag. in H.

Informatæ. Sub pede & armo dext. 16.20 H 17.30 A 4  
Trium supra australe cornu præced. 11.0 H 2.0 5  
Duarum sub extremo austr. cornu bor. 20.0 II 6.0 5  
Australina Schreckenf. 20' 20.0 II 7.0 A 5  
Ex 5 sub cornu bor. pr. Schr. 3.0' 18.0 II 0.40 B 5  
Hanc sequens 20.0 II 1.20 5  
Hac iterum posterior 22.0 II 1.40 5  
Ultimarum duarum borea 23.20 II 3.40 5  
Australis 24.20 II 1.35 B

In GEMINIS, Quæ præcedit genu anterius, clara 27.45 II 6.10 B 4  
In CANCRO. Supra flexum Forcipis australina 11.20 H 2.20 A 4  
Quæ sequitur extremâ Forc. austr. 13.20 H 5.20 A 4  
In VIRGINE. Infem. □ præced. 18.0 H 0.10 B 6  
In coxa extremo [australior] 19.0 H 8.30 B 5

Ex lib. meo de stella Nova Serpentarij.  
In LIBRA. Infra Chelâ austr. bor. 23.2 M 8.42 A 4  
Australior 23.35 M 10.20 A 4  
In SCORPIONE. In corpore, sub v. vola pedis Serpent. 2.57 H 1.50 A 4